



## Accident & Emergency Department

### Limping Child

#### Information for patients, parents and carers

#### Why is my child limping?

We see many children who have been brought to the Emergency Department because they have developed a limp (and/or pain somewhere in the affected leg). Most of these children will have an "irritable hip" (see below) or another condition which does not need investigation to diagnose and will get better on its own without hospital treatment.

There are some other more serious causes of a limp, such as joint or bone infection which do need hospital treatment. The Emergency Department Doctor will have ruled out these more serious causes by taking a history and examining your child carefully. Occasionally an x-ray will have been taken.

#### What is an Irritable Hip?

Irritable hip (or transient synovitis) is the commonest cause of limp in childhood. It is due to inflammation (**not** infection) of the hip joint. The exact cause is not known but it often happens after a viral illness or an injury.

Your child may complain of pain in the hip, groin, thigh or knee and may have difficulty walking or crawling. Usually only one side is affected. It is generally a mild condition which gets better on its own after one or two weeks.

#### Does my child need to be seen again?

Yes, you should make an appointment for your child to be seen again within 48-72 hours in the Emergency Department Clinic.

#### During that time please contact the Emergency Department immediately if your child:

- **becomes unwell**
- **develops a high temperature**
- **has increasing pain**
- **is unable to put any weight at all on the affected leg**

These are symptoms that may indicate a more serious problem.

#### What can I do to help my child at home?

You should encourage your child to rest the affected leg as much as possible over the next few days. This may mean that your child does not attend school or nursery until the clinic appointment.

Give regular pain relief medicine such as, paracetamol and/or ibuprofen. Paracetamol can be given every four - six hours (no more than four times in 24 hours). Ibuprofen can be given every six -eight hours (no more than three times in 24 hours). Do not exceed the prescribed dose. Most children will show some improvement over the next few days.

### **What will happen at the 48-72 hour appointment?**

This appointment gives us a chance to check on your child's progress. We would expect most children to be getting better at this time. Your child will be seen and examined by a senior Emergency Department Doctor.

If your child is well and the limp/pain is improving, your child will be discharged from the clinic.

If your child is unwell, not improved at all or is worse, then the doctor will probably arrange investigations e.g. blood tests or x-rays. Your child may then be referred to another team.

### **When can I expect my child to be back to normal?**

You can allow your child to gradually go back to their usual activities as they improve, but we advise they do not do sport or strenuous activity for two weeks. Your child may still need pain relief medicine. Gradually reduce the number of times you give it in a day as they get better.

We would expect your child to be back to normal within one - two weeks.

### **Please contact the Emergency Department if your child unexpectedly becomes worse in this time or does not get completely better in two weeks.**

Alder Hey Emergency Department telephone number: 0151 293 3623

This leaflet only gives general information. You must always discuss the individual treatment of your child with the appropriate member of staff. Do not rely on this leaflet alone for information about your child's treatment. This information can be made available in other languages and formats if requested.

Alder Hey Children's NHS Foundation Trust  
Eaton Road  
Liverpool  
L12 2AP

Tel: 0151 228 4811  
[www.alderhey.nhs.uk](http://www.alderhey.nhs.uk)

LIMPCA&E/F/1/10 ©Alder Hey 2015 Date of next review: March 2018 PIAG 52